

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
30 September 2004 (30.09.2004)

PCT

(10) International Publication Number
WO 2004/083214 A1

(51) International Patent Classification⁷: **C07D 491/22**,
A61K 31/437, A61P 35/00 // (C07D 491/22, 311:00,
211:00, 221:00, 209:00)

Franco [IT/IT]; c/o Istituto Nazionale per lo Studio e la
Cura dei Tumori, Via Venezian, 1, I-20133 Milan (IT).

(21) International Application Number:
PCT/IT2004/000118

(74) Agent: **SPADARO, Marco et al.**; Cavattoni-Raimondi,
Viale dei Parioli, 160, I-00197 Roma (IT).

(22) International Filing Date: 10 March 2004 (10.03.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
10/388,587 17 March 2003 (17.03.2003) US

(81) Designated States (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(71) Applicants (*for all designated States except US*):
**SIGMA-TAU INDUSTRIE FARMACEUTICHE RI-
UNITE S.p.A.** [IT/IT]; Viale Shakespeare, 47, I-00144
Roma (IT). **ISTITUTO NAZIONALE PER LO STU-
DIO E LA CURA DEI TUMORI** [IT/IT]; Via Venezian,
1, I-20133 Milano (IT).

(84) Designated States (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), Euro-
pean (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR,
GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).

(72) Inventors; and


(75) Inventors/Applicants (*for US only*): **DALLAVALLE**,
Sabrina [IT/IT]; Via Montenero, 9, I-20059 Vimercate
(IT). **PENCO, Sergio** [IT/IT]; Via Milly Carla Mignone,
5, I-20133 Milan (IT). **PISANO, Claudio** [IT/IT]; c/o
Sigma-Tau Industrie Farmaceutiche Riunite S.p., A., Via
Pontina, Km. 30,400, I-00040 Pomezia (IT). **ZUNINO**,

Published:

— with international search report

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: 7-IMINO DERIVATIVES OF CAMPTOTHECIN HAVING ANTITUMOR ACTIVITY



(I)

(57) Abstract: Camptothecin derivatives of camptothecin of formula (I), wherein the groups R₁, R₂ and R₃ are as defined in the description are disclosed. The compounds of formula (I) are endowed with antitumor activity and show a good therapeutic index. Processes for the preparation of the compounds of formula (I) and their use in the preparation of medicaments useful in the treatment of tumors, viral infections and antiparasmodium falciparum are also disclosed.

WO 2004/083214 A1

10/564637

1

IAP20 Rec'd PCT/PTO 13 JAN 2006

7-Imino derivatives of camptothecin having antitumor activity

The present invention relates to compounds having antitumor activity, in particular to new derivatives of camptothecins, processes for their preparation, their use as antitumor drugs and pharmaceutical compositions containing them as active ingredients.

Background of the invention

Camptothecin is an alkaloid, which was isolated by Wall et al. (*J. Am. Chem. Soc.* 88, 3888-3890 (1966)) for the first time from the tree *Camptoteca acuminata*, a plant originating from China, of the Nyssaceae family.

The molecule consists of a pentacyclic structure having a lactone in the ring E, which is essential for cytotoxicity.

The drug demonstrated a wide spectrum of antitumor activity, in particular against colon tumors, other solid tumors and leukemias, and the first clinical trials were performed in the early 70's. Since Camptothecin (in the following briefly CPT) has low water solubility and in order to prepare clinical trials, the National Cancer Institute (NCI) prepared the sodium salt (NSC100880), which is water-soluble. Clinical trials in phase I and II, were not completed because of the high toxicity showed by the compound (hemorrhagic cystitis, gastrointestinal toxicity, such as nausea, vomit, diarrhoea, and myelosuppression, especially leucopenia and thrombocytopenia).

In any case, sodium salt showed a lower activity than CPT, because, at pH 7.4, the inactive form (open ring) predominates on the lactone-active one (closed ring), which predominates at pH < 4.0.

Subsequently, many CPT analogues were synthesised in order to obtain compounds with lower toxicity and higher water solubility. Two drugs are marketed, Irinotecan (CPT-11), marketed with the Trade

Mark Camptosar® by Upjohn and Topotecan, marketed with the Trade Mark Hymcamptamin® or Thycantin®, by Smith Kline & Beecham. Other derivatives are in different steps of clinical development in phase II, such as NSC-603071 (9-amino-camptothecin), 9-NC or 9-nitrocampthothecin, an oral prodrug converted in 9-aminocampthothecin, GG-211 (GI 147211), and DX-8591f, the latter being water-soluble. All the derivatives identified to-date contain the parent structure with 5 rings, essential for cytotoxicity. It was demonstrated that modifications on the first ring, such as in the case of the above-mentioned drugs increase water solubility and allow a higher tolerability of the drug.

Water-soluble Irinotecan was approved for the treatment of many solid tumors and ascites (colon-rectum, skin, stomach, breast, small and non-small cell lung, cervix and ovarian cancer and in non-Hodgkin lymphoma). Moreover, Irinotecan resulted active in solid tumors resistant to Topotecan, vincristine or melphalan and MDR-1 cells resulted marginally resistant to the drug. The active metabolite was identified as the 10-hydroxyderivative (SN-38), produced by the action of carboxylesterases. CPT-11 showed a good activity using different administration routes, such as intraperitoneal, intravenous, oral (*Costin, D., Potmhexyl, M.. Advances in Pharmacol., 29B, 51-72 1994*).

CPT-11 was administered also with cisplatin or etoposide, showing a synergistic effect, thanks to the ability to hinder DNA repair. Also in this case, however, a grade 3 and 4 leucopenia and diarrhoea arose (*Sinha, B.K., (1995), Topoisomerase inhibitors. Drugs, 49, 11-19, 1995*).

Topotecan has a significant oral bioavailability. Oral administration proved to be convenient to reach a prolonged exposition to the drug, without the use of temporary catheters being necessary (*Rothenberg, M.L., Annals of Oncology, 8, 837-855, 1997*). Also this water-soluble CPT analogue showed activity against different types of tumors, with different administration routes, intraperitoneal, intravenous, subcutaneous, oral. The more promising results were obtained with


Topotecan hydrochloride, intravenous infusion for 5 days, in different tumors such as small and non-small cell lung, ovarian, breast, stomach, liver, prostate, soft tissue sarcoma, head and neck, oesophagus, resistant colon-rectum, multiform glioblastoma, chronic and acute myelocytic leukemias. However, also in this case, severe side effects occurred, such as neutropenia and thrombocytopenia, whereas gastrointestinal toxicity, such as nausea, vomit and diarrhoea were milder.

It was demonstrated that the main transformation and elimination pathways of the drug comprise lactone hydrolysis and urinary excretion: in fact, lactone form is 50% hydrolysed to open ring, 30 minutes after infusion. Topotecan crosses hematoencephalic barrier 10 minutes after infusion (30% in the cerebrospinal fluid with respect to plasma). On the contrary, camptothecin does not cross hematoencephalic barrier in significant amount, probably due to its binding with proteins.

Clinical development of 9-aminocamptothecin was hampered by its scarce water solubility. Recently, a colloidal dispersion was prepared, which made possible its entry in phase II clinical trial. Prolonged exposition (from 72 hours to 21 days) appeared to be essential to demonstrate antitumor activity, because of its short half-life (*Dahut, et al., 1994*). Responses in patients suffering from not treated colon-rectum, and breast cancer and resistant lymphoma, were noticed. The activity demonstrated against Pgp-positive tumors suggested a lack of cross-resistance against resistant MDR-1 cells. Once again, bone marrow and gastrointestinal toxicity was observed.

Lurtotecan is the most water-soluble analogue, with an activity comparable to Topotecan in vitro. Two regimens were adopted: one 30-minutes infusion a day for 5 days every 3 weeks and one 72-hours infusion one time every 3 weeks. Responses in patients suffering from, neck, ovarian, breast, liver tumour were observed. Also in this case, haematic toxicity was detected.

The molecule is the following:



9-Nitrocamptothecin is an oral prodrug rapidly converted into 9-aminocamptothecin after administration. Responses were observed in patients suffering from pancreas, ovarian, and breast cancer.

Notwithstanding the major part of tumour cells is highly sensitive to topoisomerase I inhibitors, due to the high enzyme levels, some tumoral lines result to be resistant. This is due to other mechanisms, rather than the overexpression of MDR1 and MRP (multidrug resistance associated protein) genes and of their products, P (Pgp) glycoprotein and MRP protein, respectively, for which Topotecan or CPT-11 are not very good substrates, (*Kawato, Y., et al., J. Pharm. Pharmacol., 45, 444-448, (1993)*).

In fact, it was observed that some resistant tumour cells contain mutant forms of topo I, accordingly the formation of the topo I-DNA complex is damaged or some cells lack in the carboxylesterase activity, necessary for converting CPT-11 in the active metabolite SN-38 and are thus resistant against this drug (Rothenberg, 1997, *ibid.*).

Within the drugs used in tumour therapy, the interest in inhibitors of topoisomerase I enzymes is attributed to the following considerations: a) efficacy against tumors naturally resistant to conventional drugs, topoisomerase II inhibitors included; b) the levels of the topo I enzyme remain elevated in all phases of the cycle; c) many tumors express high levels of the target enzyme; d) lack of recognition by the proteins

involved in the phenomenon of multi-drug resistance (Pgp or MRP) and absence of the detoxifying enzyme-mediated metabolism, associated to the glutathione-dependent system (glutathione peroxidase and glutathione S-transferase) (*Gerrits, CJH., et al., Brit. J. Cancer, 76, 952-962*).

Once potential clinical advantages of topoisomerase I inhibitors are taken into consideration, both in terms of antitumor activity, assayed on a wide range of tumors, and the poor induction of pharmacoresistance, the present research aims to identify topo I inhibitors with a lower toxicity with respect to the one demonstrated by the drugs on the market or in clinical phase. The factors determining the relative potency of camptothecin analogues include a) intrinsic activity of topoisomerase I inhibition; b) drug mean life; c) interaction with plasma proteins; d) the ratio between the circulating active form (lactone) and the non active one (carboxylate); e) drug sensitivity relative to cell outflow mediated by glycoprotein P or MRP; f) bond stability with topoisomerase I (*Rothenberg, 1997, ibid.*).

Among the main adverse effects of Irinotecan and other camptothecins derivatives, myelosuppression and gastrointestinal toxicity, such as diarrhoea and vomit, have been observed. Diarrhoea can have an early or late onset and can be a dose-limiting factor. Vomit and late diarrhoea are induced by many antitumor drugs, while early diarrhoea occurring during or immediately after infusion is almost specific for Irinotecan and some camptothecin derivatives.

Toxic effects occur mainly in the intestinal tract.

In order to reduce diarrhoea, CPT-11 was administered in some clinical trials, in combination with loperamide, a synthetic opioid, agonist of the mu-opioid enteric receptors (*Abigeres, 1994; Abigeres, 1995*), as well as with an inhibitor of the enkephalinases (acetorfan) or with ondansetron, an antagonist of the 5-HT₃ receptors, or with diphenidramine, an antagonist of H₁ receptors.

To date, the problems connected with the use of camptothecin derivatives as antitumor drugs can be summarised in the following items:

- camptothecin (CPT), and many of its active derivatives have low water solubility;
- the subsequent derivatives are endowed with severe side effects at gastrointestinal and bone marrow level;
- some tumour lines developed resistance against topoisomerase I inhibitors;
- there is the constant search for a better therapeutic index.

Patent application WO 97/31003 discloses derivatives of camptothecins substituted at positions 7, 9 and 10. Position 7 provides the following substitutions: -CN, -CH(CN)-R₄, -CH=C(CN)-R₄, -CH₂-CH=C(CN)-R₄, -C(=NOH)-NH₂, -CH=C(NO₂)-R₄, -CH(CN)-R₅, -CH(CH₂NO₂)-R₅, 5-tetrazolyl, 2-(4,5-dihydroxazolyl), 1,2,4-oxadiazolidin-3-yl-5-one, wherein R₄ is hydrogen, linear or branched alkyl from 1 to 6 carbon atoms, nitrile, carboxyalkoxy. Of these possible compounds, WO 97/31003 enables the disclosure only of camptothecin derivatives bearing at position 7 the group -CN and -CH=C(CN)₂, with unsubstituted positions 9 and 10.

Of these compounds, the best one proved to be the 7-nitrile (R₄ = -CN), hereinafter named CPT 83, with cytotoxic activity on non-small cells lung carcinoma (non-SCLC, H-460). This tumour line is intrinsically resistant to cytotoxic therapy and is only moderately responsive to topoisomerase I inhibitors, notwithstanding the overexpression of the target enzyme. CPT 83 is more active than Topotecan, taken as reference compound and on the whole it offers a better pharmacological profile, even in terms of tolerability, then a better therapeutic index.

CPT 83 is prepared through a synthesis route comprising the oxidation of 7-hydroxymethylcamptothecin to camptothecin 7-aldehyde, the

transformation of the latter into oxime and final conversion into nitrile.


The starting compound and the intermediates are disclosed in *Sawada et al., Chem. Pharm. Bull., 39, (10), 2574, (1991)*. This paper makes reference to a patent family with priority of 1981, for example European patent application EP 0 056 692, published in 1982. In these publications there are disclosed, among others, the compounds camptothecin 7-aldehyde and its oxime. The usefulness of these derivatives is to provide compounds with antitumor activity having low toxicity starting from 7-hydroxymethylcamptothecin. In the paper published on *Chem. Pharm. Bull., 39, (10) 2574, (1991)*, the authors demonstrate that, with respect to camptothecin, the 7-alkyl and 7-acyloxymethyl derivatives, which were not foreseen in the above mentioned patent application, are the more active compounds on lines of murine leukemia L1210, while lower activity, always with respect to camptothecin, was observed in compounds bearing 7-substitutions with high polar character, such as hydrazones and the oxime - CH(=NOH).

EP 1 044 977 discloses camptothecin derivatives bearing an oxime O-substituted at position 7. The general formula comprises also camptothecin derivatives bearing an enamine group at position 7. In this reference, the main teaching is directed to the antitumor activity of oxime derivatives, while imines are given in only few examples of synthetic preparation, but no pharmacological data are provided. Subsequent work of the inventors of the above mentioned patent was focused on oxime derivatives, in particular the tert-butoxy one, which, under the name of Gimatecan is now under clinical trial. Imines were considered just an alternative to oximes, but first pharmacological assays discouraged further development of this class.

Abstract of the invention

It has now surprisingly been found that camptothecins bearing an aromatic enamino group on position 7 are endowed with antitumor activity. Said compounds have better therapeutic index.

Accordingly, it is an object of the present invention compounds of general formula (I):



wherein: R_1 is a $-C(R_5)=N-R_4$ group, wherein R_4 is a phenyl group, optionally substituted with one or more groups selected from the group consisting of: halogen, hydroxy, keto, C_1 - C_8 alkyl, C_1 - C_8 alkoxy, phenyl, cyano, nitro, $-NR_6R_7$, wherein R_6 and R_7 , the same or different between them, are hydrogen, $(C_1$ - $C_8)$ linear or branched alkyl; $-S-S-(2$ -aminophenyl), $-S-S-(4$ -aminophenyl), $-S-(4$ -aminophenyl), $-SCH_3$ and $-CH_2ON=C(CH_3)_2$;

R_5 is hydrogen, C_1 - C_8 linear or branched alkyl, C_1 - C_8 linear or branched alkenyl, C_3 - C_{10} cycloalkyl, $(C_3$ - $C_{10})$ cycloalkyl - $(C_1$ - $C_8)$ linear or branched alkyl, C_6 - C_{14} aryl, $(C_6$ - $C_{14})$ aryl - $(C_1$ - $C_8)$ linear or branched alkyl;

R_2 and R_3 , the same or different between them are hydrogen, hydroxy, C_1 - C_8 linear or branched alkoxy;

their N_1 -oxides, their single isomers, in particular the syn and anti isomers of the $-C(R_5)=N-R_4$ group, their possible enantiomers,

diastereoisomers and relative admixtures, the pharmaceutically acceptable salts thereof and their active metabolites.

The present invention comprises the use of the compounds of the above-mentioned formula (I) as active ingredients for medicaments, in particular for medicaments useful for the treatment of tumors. A further object of the present invention is also the use of the compounds of formula (I) as active ingredients for medicaments useful for treating viral infections. Another object of the present invention is also the use of the compounds of formula (I) as active ingredients for medicaments having antiparasitic activity.

The present invention comprises pharmaceutical compositions containing compounds of formula (I) as active ingredients, in admixture with pharmaceutically acceptable vehicles and excipients.

The present invention comprises also processes for the preparation of compounds of formula (I), and the relative key intermediates.

Detailed description of the invention

Within the scope of the present invention, as examples of C₁-C₈ linear or branched alkyl group, methyl, ethyl, propyl, butyl, pentyl, octyl are meant and their possible isomers, such as for example isopropyl, isobutyl, tert-butyl.

As halogen it is intended fluorine, chlorine, bromine, iodine.

Examples of pharmaceutically acceptable salts are, in case of nitrogen atoms having basic character, the salts with pharmaceutically acceptable acids, both inorganic and organic, such as for example, hydrochloric acid, sulfuric acid, acetic acid, or, in the case of acid group, such as carboxyl, the salts with pharmaceutically acceptable bases, both inorganic and organic, such as for example, alkaline and

alkaline-earth hydroxides, ammonium hydroxide, amine, also heterocyclic ones.


A first group of preferred compounds is the one wherein R_4 is phenyl substituted by at least one a residue selected from the group consisting of methyl, ter-butyl, methoxy, hydroxy, chloro, iodo, nitro, $-S-S-(2\text{-aminophenyl})$, $-S-S-(4\text{-aminophenyl})$, $-S-(4\text{-aminophenyl})$, $-SCH_3$ and $-CH_2ON=C(CH_3)_2$. In particular, the phenyl group is most preferably substituted in ortho-position.

A second group of particularly preferred compounds comprises:

7-(2-methylphenyl)iminomethylcamptothecin (ST2212)
7-(2-chlorophenyl)iminomethylcamptothecin (ST2228)
7-(2,6-dimethylphenyl)iminomethylcamptothecin (ST2317)
7-(2-iodophenyl)iminomethylcamptothecin (ST2316)
7-(2-methoxyphenyl)iminomethylcamptothecin (ST2343)
7-(4-methylphenyl)iminomethylcamptothecin (ST2478)
7-(2-hydroxyphenyl)iminomethylcamptothecin (ST2389)
7-(4-chlorophenyl)iminomethylcamptothecin (ST2412)
7-(4-methoxyphenyl)iminomethylcamptothecin (ST2477)
7-[(4-isopropylidene-amino-oxymethyl)phenyl]iminomethylcamptothecin (ST2460)
7-(2-t-butylphenyl)iminomethylcamptothecin (ST2388)
7-phenyliminomethylcamptothecin (ST1546)
7-(4-nitrophenyl)iminomethylcamptothecin (ST1561)
7-2-(2-aminophenyldithio)phenyliminomethylcamptothecin (ST1737)
7-4-(4-aminophenyldithio)phenyliminomethylcamptothecin (ST2034)
7-4-(4-aminophenylthio)phenyliminomethylcamptothecin (ST2069)
7-(2-methylthiophenyl)iminomethylcamptothecin (ST2138)
7-(4-tert-butylphenyliminomethyl)-camptothecin (ST 2619)
7-(4-methylthiophenyliminomethyl)-camptothecin (ST 2667)
7-(4-hydroxyphenyliminomethyl)-camptothecin (ST 2616)

The compounds of formula (I) can be prepared with different methods according to the nature of the R_4 group linked to the nitrogen of the 7-iminomethyl group.

The compounds of formula (I) wherein R_4 is as above defined can be prepared starting from camptothecin 7-aldehyde (formula Ia, R_5 hydrogen) or 7-keto camptothecin (formula Ia, R_5 different from hydrogen),



wherein R_1 is the group $-C(R_5)=O$, and R_5 is as defined for the formula (I), R_2 and R_3 are as defined in formula (I). The compound of formula (Ia) is reacted with the compound of formula (IIb) R_4-NH_2 , wherein R_4 is as above, to give compounds of formula (I), wherein R_1 is the group $-C(R_5)=N-R_4$, R_4 is defined as in formula 1. The reaction can be carried out with conventional methods well known to the person skilled in the art, being a normal formation of an imine. Preferably, the molar ratio between 7-aldehyde or 7-keto camptothecin and amine is comprised between 1:3 and 3:1. The salts of the amine of interest can also be used. The reaction is carried out in the presence of a base, for example an inorganic base, such as potassium carbonate, or organic, such as triethylamine or diazabicyclononene, using polar solvents, preferably methanol or ethanol and carrying out the reaction at a temperature comprised between room temperature and solvent boiling point, optionally in the presence of dehydrating agents, for example sodium or magnesium sulfate, molecular sieves. If desired it is also possible to carry out the reaction in the presence of a catalyst, for example a

Lewis acid (as disclosed for example by *Moretti and Torre, Synthesis, 1970, 141*; or by *Kobayashi, et al., Synlett, 1977, 115*).

The camptothecin 7-aldehyde and the camptothecin 7-oxime are disclosed in the patent application EP 0 056 692 and in the mentioned *Sawada, et al., Chem. Pharm. Bull., 39, (10) 2574 (1991)*.

N₁-oxides of the compounds of formula (I) are prepared according to well-known methods of oxidation of heteroaromatic nitrogen, preferably by oxidation with acetic or trifluoroacetic acid and hydrogen peroxide, or by reaction with organic peroxyacids (*A. Albini and S. Pietra, Heterocyclic N-oxides, CRC, 1991*).

Regarding the various meanings of R₄, present in the different reactivities of formula II, these reactivities are available in the market, or can be prepared according to well-known methods in literature, which the expert in the field can resort to, completing with their own knowledge of the argument.

Pharmaceutically acceptable salts are obtained with conventional methods found in the literature, and do not necessitate of further disclosure.

The compounds disclosed in the present invention show antiproliferative activity, therefore are useful for their therapeutical activity, and possess physico-chemical properties that make them suitable to be formulated in pharmaceutical compositions.

The pharmaceutical compositions comprise at least a compound of formula (I), in an amount such as to produce a significant therapeutical effect, in particular antitumoral effect. The compositions comprised within the present invention are conventional and are obtained with commonly used methods in the pharmaceutical industry. According to the desired administration route, the compositions shall be in solid or liquid form, suitable to the oral, parenteral, intravenous

route. The compositions according to the present invention comprise together with the active ingredients at least a pharmaceutically acceptable vehicle or excipient. Formulation co-adjuvants, for example solubilizing, dispersing, suspending, emulsifying agents can be particularly useful.

The compounds of formula (I) can also be used in combination with other active ingredients, for example other antitumor drugs, both in separate forms, and in a single dose form.

The compounds according to the present invention are useful as medicaments with antitumor activity, for example in lung tumors, such as the non-small cell lung tumour, tumors of the colon-rectum, prostate, gliomas.

Cytotoxic activity of the compounds of the present invention was assayed in cell systems of human tumour cells, using the antiproliferative activity test as a method of evaluation of the cytotoxic potential.

The cell line used is a lung non-small cell carcinoma that belongs to non-small cells hystotype named NCI H460.

For the *in vivo* studies, the solubilization was carried out in 10% DMSO in bidistilled water, being impossible the solubilization in saline, and the administration for the oral route was carried out at a volume of 10 ml/kg.

Antitumoral activity

Atimic nu/nu Swiss mice (Charles River, Calco, Italia), ageing 10-12 weeks were used. The animals were maintained in laminar flow rooms, according to the guidelines of the United Kingdom Co-ordination Committee Cancer Research. Experimental protocols were approved by

the Ethical Committee for animal experimentation of Istituto Nazionale per lo Studio e la Cura dei Tumori.

Tumour fragments of about 2x2x2 mm coming from mice to which were inoculated s.c. 10^6 cells NCI H460/mouse, were implanted s.c. bilaterally in groups of 5 mice each.

The animals were treated with the compounds when the tumour began to be palpable.

Twice a week, using a Vernier calliper, the width, minimum diameter (l), length and maximum diameter (L) of the tumors were measured, in mm. The tumour volume (mm^3) was calculated according to the formula $l^2 \times L / 2$. Efficacy of the molecule was evaluated as TVI percent of the treated group versus the control group according to the formula $\text{TVI}\% = 100 - (T/C \times 100)$, wherein T is the mean value of the tumour volume of the treated group and C of control one. A compound is considered active when $\text{TVI}\% \geq 50$.

Further advantages of these molecules can be identified in the wide interval of effective doses, indicating an increase of therapeutic index and a higher handling in the therapeutical use, in particular if a prolonged administration in the time is foreseen, above all in the injectable formulations, with the use of variable schemes and doses.

An important drawback of conventional camptothecins is the reversibility of their bond in the ternary complex (drug-DNA-enzyme). This reversibility affects drug efficacy, as it does not allow the transformation of the single strand DNA cleavage into double strand DNA cleavage during DNA synthesis.

The advantage offered by the compounds according to the present invention is evident in overcoming the limit of reversibility of the ternary complex with respect to the state of the art.

In preclinical investigations, the compounds of the present invention showed cytotoxic activity in various tumor cell lines.

This broad spectrum of anticancer activity was confirmed in mice transplanted with human tumor xenografts, including NSCLC (H460, A549), prostate ca. (JCA-1), glioblastoma (GBM/7), gastric cancer. (MKN28), osteosarcoma (U2OS), ovarian cancer (A2780/Dx, A2780/DDP) and colon (HT29, CoBA) carcinomas as well as in murine lung cancer (M109) and leukaemia model (L1210).

The preclinical data suggest that the compounds of the present invention may be an active anticancer agent against human's cancers and in particular against non-small cell lung cancer (NSCLC), glioblastoma and prostate carcinoma.

The antitumor activity of exemplary compounds of the invention is shown in the following Table 1.

Table 1

Compound	IC ₅₀ (mg/ml)	
ST 1737	0.0295±0.07	(water soluble)
ST 2034	0.235±0.02	
ST 2069	0.14±0.04	
ST 2138	0.033±0.004	
ST 2228	0.036±0.01	
ST 2316	0.062±0.03	
ST 2317	0.075±0.02	
ST 1561 (CPT160)	0.14±0.03	
ST 2412	0.042±0.0012	
ST 2212	0.235±0.06	
ST 2388	0.033±0.004	
ST 2389	0.015±0.008	
ST 2343	0.028±0.012	
ST 2477	0.08±0.002	

The in vivo activity of the compound ST1737 is shown in Table 2 below.

Table 2

Effects of ST 1737 administered per os, q4dx4, in athymic nude mice bearing s.c. the MKN-28 human gastric carcinoma

Drug	Dose (mg/kg)	TVI% ¹	BWL% ²	Tox ³
ST 1737	4	69		0/4
	7	56		0/4
	12	68	4	0/4

Mean tumor doubling times in control mice were 7.1±1.3 and 5.4±1.4 days (exp. 656 and 664, respectively).

¹Tumor volume inhibition % in treated over control tumors at 11-12 day after last treatment.

²Body weight loss % induced by drug treatment.

³Dead/treated mice.

The high cytotoxic potency of the compounds of the present invention, herein represented in an exemplary way with one of the preferred compounds, ST 1737, is also reflected by the potent antitumor activity. Using a panel of tumor xenografts characterized by a significant responsiveness to Topotecan (TPT) (i.e. TVI > 80%), the spectrum of antitumor activity of the compounds of the present invention, against a significant number of human tumor models was substantially improved. In particular, an impressive antitumor efficacy was found in the treatment of many tumor models, where very high regressions were achieved in a large number of treated animals. Moreover, the compounds of the present invention, were able to induce substantial CR in the tumors characterized by a MDR-phenotype. This observation is of high importance, indicating that the compounds of the present invention are not a substrate for P-glycoprotein.

Additional therapeutic advantages of the compounds of the present invention are related to a) an improvement of the therapeutic index, b) drug efficacy in a large range of doses, c) evidence of efficacy using quite different schedules, making the compounds of the present invention less dependent on the treatment schedule than that of Topotecan.

The following examples further illustrate the invention.

Example 1

General procedure: To a suspension of Yb(OTf)₃ (16 mg, 0.03 mmol) in 5 ml of anhydrous CH₂Cl₂ containing 4 Å MS, a solution of 7-formylcamptothecin (100 mg, 0.26 mmol) in 20 ml of CH₂Cl₂ is added, followed by a solution of the amine (0.26 mmol) in 0.5 ml of CH₂Cl₂. The resulting mixture is stirred at room temperature until the reaction is complete. After filtering the sieves, 20 ml of water are added and the two phases are separated. The aqueous layer is rapidly extracted three times with dichloromethane. The combined organic phases are dried

and evaporated, and the product purified by flash chromatography on silica gel.

The following compounds were obtained. In some cases antitumor activity is shown (IC₅₀ on H-460, μ M)

7-(2-methylphenyl)iminomethylcamptothecin (ST2212)

IC₅₀ (H-460, μ M): 0.10

M.P. 247-248 °C dec., ¹H NMR (DMSO-d₆) δ : 0.87 (t, J = 7 Hz, H₃-18), 1.7-1.9 (m, H₂-19), 2.5 (s, Ar-CH₃), 5.4 (s, H₂-17), 5.60 (s, H₂-5), 6.55 (s, -OH), 7.25-7.50 (m, 4H Ar, H-14), 7.75 (m, H-11), 7.95 (m, H-10), 8.25 (dd, H-12), 9.10 (dd, H-9), 9.65 (s, CH=N).

7-(2-chlorophenyl)iminomethylcamptothecin (ST2228)

IC₅₀ (H-460, μ M): 0.07

M.P. >240°C dec., ¹H NMR (DMSO-d₆) δ : 0.83 (t, J = 7 Hz, H₃-18), 1.7-1.9 (m, H₂-19), 5.45 (s, H₂-17), 5.60 (s, H₂-5), 6.50 (s, -OH), 7.35-7.50 (m, H-14; 4H Arom.), 7.85 (m, H-11), 7.95 (m, H-10), 8.30 (dd, H-12), 9.10 (dd, H-9), 9.70 (s, CH=N).

7-(2,6-dimethylphenyl)iminomethylcamptothecin (ST2317)

IC₅₀ (H-460, μ M): 0.15

M.P. 250 °C dec., ¹H NMR (DMSO-d₆) δ : 0.87(t, J = 7 Hz, H₃-18), 1.7-1.9 (m, H₂-19), 2.25 (s, 2Ar-CH₃), 5.4 (s, H₂-17), 5.60 (s, H₂-5), 6.55 (s, -OH), 7.0-7.30 (m, 3H Ar), 7.40 (s, H-14), 7.8 (m, H-11), 7.9 (m, H-10), 8.25 (dd, H-12), 8.85 (dd, H-9), 9.5 (s, CH=N).

7-(2-iodophenyl)iminomethylcamptothecin (ST2316)

IC₅₀ (H-460, μ M): 0.06

M.P. 240 °C dec., ¹H NMR (DMSO-d₆) δ : 0.87(t, J = 7 Hz, H₃-18), 1.8-1.9 (m, H₂-19), 5.45 (s, H₂-17), 5.75 (s, H₂-5), 6.55 (s, -OH), 7.1-7.6 (m, 4H Ar, H-14), 7.8 (m, H-11), 7.9 (m, H-10), 8.30 (dd, H-12), 9.10 (dd, H-9), 9.65 (s, CH=N).

7-(2-methoxyphenyl)iminomethylcamptothecin (ST2343)

IC₅₀ (H-460, μ M): 0.06

M.P. 244-246°C dec., ¹H NMR (DMSO-d₆) δ : 0.83 (t, J = 7 Hz, H₃-18), 1.7-1.9 (m, H₂-19), 3.95 (s, OCH₃), 5.45 (s, H₂-17), 5.55 (s, H₂-5), 6.45 (s, -OH), 7.0-7.50 (m, H-14; 4H Arom.), 7.7 (m, H-11), 7.85 (m, H-10), 8.25 (dd, H-12), 8.9 (dd, H-9), 9.70 (s, CH=N).

7-(4-methylphenyl)iminomethylcamptothecin (ST2478)

IC₅₀ (H-460, μ M): 0.18

M.P. 159-160°C dec., ¹H NMR (DMSO-d₆) δ : 0.83 (t, J = 7 Hz, H₃-18), 1.7-1.9 (m, H₂-19), 2.35 (s, Ar-CH₃), 5.37 (s, H₂-17), 5.5 (s, H₂-5), 6.45 (s, -OH), 7.25-7.35 (m, H-14; 2H Arom.), 7.4-7.5 (m, 2H arom.), 7.7 (m, H-11), 7.85 (m, H-10), 8.16 (dd, H-12), 8.9 (dd, H-9), 9.55 (s, CH).

7-(2-hydroxyphenyl)iminomethylcamptothecin (ST2389)

IC₅₀ (H-460, μ M): 0.06

M.P. 252-254 °C dec., ¹H NMR (DMSO-d₆) δ : 0.87 (t, J = 7 Hz, H₃-18), 1.7-1.9 (m, H₂-19), 5.4 (s, H₂-17), 5.60 (s, H₂-5), 6.55 (s, -OH), 6.90-7.5 (m, 4H Ar, H-14), 7.85-8.0 (m, H-11, H-10), 8.35 (dd, H-12), 8.90 (dd, H-9), 9.70 (s, CH=N).

7-(4-chlorophenyl)iminomethylcamptothecin (ST2412)

IC₅₀ (H-460, μ M): 0.08

M.P. 246-247°C dec., ¹H NMR (DMSO-d₆) δ : 0.83 (t, J = 7 Hz, H₃-18), 1.7-1.9 (m, H₂-19), 5.40 (s, H₂-17), 5.55 (s, H₂-5), 6.45 (s, -OH), 7.35 (s, H-14), 7.50-7.60 (m, 4H arom.), 7.85 (m, H-11), 7.95 (m, H-10), 8.25 (dd, H-12), 8.95 (dd, H-9), 9.55 (s, CH=N).

7-(4-methoxyphenyl)iminomethylcamptothecin (ST2477)

IC₅₀ (H-460, μ M): 0.16

M.P. 252-255 °C dec., ¹H NMR (DMSO-d₆) δ : 0.87(t, J = 7 Hz, H₃-18), 1.7-1.9 (m, H₂-19), 3.8 (s, -OCH₃), 5.4 (s, H₂-17), 5.45 (s, H₂-5), 6.55 (s, -OH), 7.05 (d, 2H Ar), 7.35 (s, H-14), 7.60 (d, 2H Ar), 7.85 (m, H-11), 7.9 (m, H-10), 8.25 (dd, H-12), 8.8 (dd, H-9), 9.5 (s, CH).

7-[(4-isopropylidene-amino-oxymethyl)phenyl]iminomethylcamptothecin (ST2460)

IC₅₀ (H-460, μ M): 0.01

M.P. 147°C dec., ¹H NMR (DMSO-d₆) δ : 0.85 (t, J = 7 Hz, H₃-18), 1.7-1.9 (m, H₂-19, C(CH₃)₂), 5.05 (s, CH₂-O), 5.40-5.55 (m, H₂-17, H₂-5), 6.50 (s, -OH), 7.35 (s, H-14), 7.40-7.60 (m, 4H arom.), 7.75-7.85 (m, H-11), 7.86-7.95 (m, H-10), 8.25 (dd, H-12), 8.95 (dd, H-9), 9.60 (s, CH=N).

7-(2-t-butylphenyl)iminomethylcamptothecin (ST2388)

IC₅₀ (H-460, μ M): 0.07

M.P. 215°C dec., ¹H NMR (DMSO-d₆) δ : 0.85 (t, J = 7 Hz, H₃-18), 1.45 (s, 9H, tBut), 1.7-1.9 (m, H₂-19), 5.35-5.75 (m, H₂-17, H₂-5), 6.50 (s, -OH), 7.05-7.5 (m, H-14; 4H arom.), 7.75-7.85 (m, H-11), 7.88-7.95 (m, H-10), 8.25 (dd, H-12), 8.95 (dd, H-9), 9.45 (s, CH=N).

7-phenyliminomethylcamptothecin (ST1546)

IC₅₀ (H-460, μ M): 0.13

¹H NMR (DMSO-d₆) δ : 0.83 (t, J = 7 Hz, H₃-18), 1.7-1.9 (m, H₂-19), 5.37 (s, H₂-17), 5.5 (s, H₂-5), 6.45 (s, -OH), 7.25-7.35 (m, H-14; H arom.), 7.4-7.5 (m, 4H arom), 7.7 (m, H-11), 7.85 (m, H-10), 8.16 (dd, H-12), 8.9 (dd, H-9), 9.55 (s, CH=N)

7-(4-nitrophenyl)iminomethylcamptothecin (ST1561)

IC₅₀ (H-460, μ M): 0.28

M.P. 260-265°C dec. ¹H NMR (DMSO-d₆) δ : 0.85 (t, J = 7 Hz, H₃-18), 1.7-1.9 (m, H₂-19), 5.35 (s, H₂-17), 5.48 (s, H₂-5), 6.45 (s, -OH), 7.3 (s, H-14), 7.6-7.7 (m, 2 Ar), 7.8 (m, H-11), 7.9 (m, H-10), 8.25 (dd, H-12), 8.35-8.40 (m, 2 Ar), 8.9 (dd, H-9), 9.67 (s, CH=N).

7-2-(2-aminophenyldithio)phenyliminomethylcamptothecin (ST1737)

IC₅₀ (H-460, μ M): 0.017

¹H NMR (DMSO-d₆) δ : 0.83 (t, J = 7 Hz, H₃-18) 1.7-1.9 (m, H₂-19) 5.35-5.75 (6H, m, H₂-5 + H-17 + NH₂), 6.40 (1H, m, ArH), 6.5-6.6 (2H, m, 1 ArH + OH), 6.90 (1H, m, ArH), 7.25-7.45 (4H, m, 3 ArH + H-14), 7.15-8.0 (4H, m, 4 ArH), 8.25 (1H, dd), 9.75 (1H, s, CH=N).

7-4-(4-aminophenyldithio)phenyliminomethylcamptothecin (ST2034)

IC₅₀ (H-460, μ M): 0.39

M.P. 154-155°C dec., ¹H NMR (DMSO-d₆) δ : 0.83 (t, J = 7 Hz, H₃-18), 1.7-1.9 (m, H₂-19), 5.40 (s, H₂-17), 5.55 (s, H₂-5 + NH₂), 6.50 (s, -OH),

6.55 (m, 2H Ar), 7.25 (m, 2H Ar), 7.35 (s, H-14), 7.60 (m, 4H Ar), 7.8 (m, H-11), 7.9 (m, H-10), 8.25 (dd, H-12), 9.0 (dd, H-9), 9.70 (s, CH=N).

7-4-(4-aminophenylthio)phenyliminomethylcamptothecin (ST2069)

IC₅₀ (H-460, μ M): 0.24

M.P. 187-188°C dec., ¹H NMR (DMSO-d₆) δ : 0.87 (t, J = 7 Hz, H₃-18), 1.7-1.9 (m, H₂-19), 5.4 (s, H₂-17), 5.55 (s, H₂-5 + NH₂), 6.55 (s, -OH), 6.65 (m, 2H Ar), 7.10-7.50 (m, 6H Ar + H-14), 7.8 (m, H-11), 7.9 (m, H-10), 8.30 (dd, H-12), 9.0 (dd, H-9), 9.5 (s, CH=N).

7-(2-methylthiophenyl)iminomethylcamptothecin (ST2138)

IC₅₀ (H-460, μ M): 0.06

M.P. >250°C dec., ¹H NMR (DMSO-d₆) δ : 0.83 (t, J = 7 Hz, H₃-18), 1.7-1.9 (m, H₂-19), 2.50 (s, SCH₃), 5.40 (s, H₂-17), 5.70 (s, H₂-5), 6.45 (s, -OH), 7.25-7.35 (m, H-14; 3H arom.), 7.6 (m, 1H arom), 7.8 (m, H-11), 7.95 (m, H-10), 8.30 (dd, H-12), 9.10 (dd, H-9), 9.55 (s, CH=N).

Example 2

To a suspension of 20S-camptothecin-7-aldehyde (1) (100 mg, 0.26 mmol) in 7 mL anhydrous CH₂Cl₂, the appropriate amine (0.78 mmol) and Yb(OTf)₃ (16 mg, 0.03 mmol) were added. The resulting mixture was stirred at room temperature until the reaction was complete. After filtering the sieves the solvent was evaporated, and the product purified by flash chromatography on silica gel (Merck 230-400 mesh).

The following compounds were obtained

7-(4-tert-butylphenyliminomethyl)-camptothecin (ST 2619)

The solution is stirred 1.5 h. Flash chromatography (eluent: CH₂Cl₂:MeOH 99:1). Yellow powder. Yield 50%, M.P. 250°C dec., ¹H

NMR (DMSO-d₆) δ : 0.88 (t, J = 7 Hz, H3-18), 1.30 (s, tBut), 1.75-1.95 (m, H2-19), 5.45 (s, H2-17), 5.55 (s, H2-5), 6.55 (s, -OH), 7.35 (s, H-14), 7.45-7.60 (m, 4H Ar), 7.80 (m, H-11), 7.95 (m, H-10), 8.25 (dd, H-12), 8.95 (dd, H-9), 9.7 (s, CH=N).

IC₅₀ (H-460, μ M): 0.09

7-(4-methylthiophenyliminomethyl)-camptothecin (ST 2667)

The solution is stirred 22h. Flash chromatography (eluent: CH₂Cl₂:MeOH 98 : 2). Yellow powder. Yield 36%, M.P. 160°C dec., ¹H NMR (DMSO-d₆) δ : 0.87 (t, J = 7 Hz, H3-18), 1.7-1.9 (m, H2-19), 2.55 (s, -SCH₃), 5.45 (s, H2-17), 5.55 (s, H2-5), 6.50 (s, -OH), 7.35 (s, H-14), 7.40 (d, 2H Ar), 7.55 (d, 2H Ar), 7.80 (m, H-11), 7.9 (m, H-10), 8.20 (dd, H-12), 8.95 (dd, H-9), 9.7 (s, CH=N).

IC₅₀ (H-460, μ M): 0.074


7-(4-hydroxyphenyliminomethyl)-camptothecin (ST 2616)

The solution is stirred 3h. Flash chromatography (eluent: CH₂Cl₂:MeOH 96:4). Yellow powder. Yield 79%, M.P. 250 °C dec., ¹H NMR (DMSO-d₆) δ : 0.90 (t, J = 7 Hz, H3-18), 1.75-2.0 (m, H2-19), 5.4 (s, H2-17), 5.55 (s, H2-5), 6.50 (s, -OH), 6.90 (d, 2H Ar), 7.35 (s, H-14), 7.55 (d, 2H Ar), 7.80 (m, H-11), 7.90 (m, H-10), 8.25 (dd, H-12), 9.0 (dd, H-9), 9.70 (s, CH=N).

IC₅₀ (H-460, μ M): 0.22

CLAIMS

1. Compounds of formula (I)



wherein: R_1 is a $-C(R_5)=N-R_4$ group, wherein R_4 is a phenyl group, optionally substituted with one or more groups selected from the group consisting of: halogen, hydroxy, keto, C_1 - C_8 alkyl, C_1 - C_8 alkoxy, phenyl, cyano, nitro, $-NR_6R_7$, wherein R_6 and R_7 , the same or different between them, are hydrogen, $(C_1$ - $C_8)$ linear or branched alkyl; $-S-S-(2$ -aminophenyl), $-S-S-(4$ -aminophenyl), $-S-(4$ -aminophenyl), $-SCH_3$ and $-CH_2ON=C(CH_3)_2$;

R_5 is hydrogen, C_1 - C_8 linear or branched alkyl, C_1 - C_8 linear or branched alkenyl, C_3 - C_{10} cycloalkyl, $(C_3$ - $C_{10})$ cycloalkyl - $(C_1$ - $C_8)$ linear or branched alkyl, C_6 - C_{14} aryl, $(C_6$ - $C_{14})$ aryl - $(C_1$ - $C_8)$ linear or branched alkyl;

R_2 and R_3 , the same or different between them are hydrogen, hydroxy, C_1 - C_8 linear or branched alkoxy;

their N_1 -oxides, their single isomers, in particular the syn and anti isomers of the $-C(R_5)=N-R_4$ group, their possible enantiomers, diastereoisomers and relative admixtures, the pharmaceutically acceptable salts thereof and their active metabolites.

2. Compounds according to claim 1, wherein R_4 is phenyl substituted by at least one a residue selected from the group consisting of methyl, ter-butyl, methoxy, hydroxy, chloro, iodo, nitro, $-S-S-(2\text{-aminophenyl})$, $-S-S-(4\text{-aminophenyl})$, $-S-(4\text{-aminophenyl})$, $-SCH_3$ and $-CH_2ON=C(CH_3)_2$; their N_1 -oxides, their single isomers, in particular the syn and anti isomers of the $-C(R_5)=N-R_4$ group, their possible enantiomers, diastereoisomers and relative admixtures, the pharmaceutically acceptable salts thereof and their active metabolites.


3. Compounds according to claim 1 or 2, wherein the phenyl group is substituted in ortho-position.

4. Compounds according to claim 1 or 2, selected from the group consisting of:

- 7-(2-methylphenyl)iminomethylcamptothecin (ST2212)
- 7-(2-chlorophenyl)iminomethylcamptothecin (ST2228)
- 7-(2,6-dimethylphenyl)iminomethylcamptothecin (ST2317)
- 7-(2-iodophenyl)iminomethylcamptothecin (ST2316)
- 7-(2-methoxyphenyl)iminomethylcamptothecin (ST2343)
- 7-(4-methylphenyl)iminomethylcamptothecin (ST2478)
- 7-(2-hydroxyphenyl)iminomethylcamptothecin (ST2389)
- 7-(4-chlorophenyl)iminomethylcamptothecin (ST2412)
- 7-(4-methoxyphenyl)iminomethylcamptothecin (ST2477)
- 7-[(4-isopropylidene-amino-oxymethyl)phenyl]iminomethylcamptothecin (ST2460)
- 7-(2-t-butylphenyl)iminomethylcamptothecin (ST2388)
- 7-phenyliminomethylcamptothecin (ST1546)
- 7-(4-nitrophenyl)iminomethylcamptothecin (ST1561)
- 7-2-(2-aminophenyldithio)phenyliminomethylcamptothecin (ST1737)
- 7-4-(4-aminophenyldithio)phenyliminomethylcamptothecin (ST2034)
- 7-4-(4-aminophenylthio)phenyliminomethylcamptothecin (ST2069)
- 7-(2-methylthiophenyl)iminomethylcamptothecin (ST2138)
- 7-(4-tert-butylphenyliminomethyl)-camptothecin (ST 2619)
- 7-(4-methylthiophenyliminomethyl)-camptothecin (ST 2667)
- 7-(4-hydroxyphenyliminomethyl)-camptothecin (ST 2616),

their N₁-oxides, their single isomers, in particular the syn and anti isomers of the $-C(R_5)=N-R_4$ group, their possible enantiomers, diastereoisomers and relative admixtures, the pharmaceutically acceptable salts thereof and their active metabolites.

5. A process for the preparation of the compounds of claims 1-4, comprising the reaction of a compound of formula (Ia)



wherein R₁ is the group $-C(R_5)=O$, and R₅ is as defined for the formula (I), R₂ and R₃ are as defined in formula (I), with the compound of formula (IIa) R₄-NH₂, and optional transformation of the obtained compounds of formula (I) into their N₁-oxides, their single isomers, in particular the syn and anti isomers of the $-C(R_5)=N-R_4$ group, their possible enantiomers, diastereoisomers and relative admixtures, the pharmaceutically acceptable salts thereof.

6. Process according to claim 5, wherein the molar ratio between compound of formula (Ia) and compound of formula (IIa) is comprised between 1:3 and 3:1.

7. Compounds according to any one of claims 1-4, as medicaments.

8. Pharmaceutical composition comprising a therapeutically effective amount of at least a compound of claims 1-4, in admixture with pharmaceutically acceptable vehicles and excipients.

9. Pharmaceutical composition comprising a therapeutically effective amount of at least a compound of claims 1-4, in admixture with pharmaceutically acceptable vehicles and excipients and optionally in combination with other active ingredients.
10. Pharmaceutical composition according to claim 9 wherein said other active ingredient is an antitumoral.
11. Use of a compound of claims 1-4, for the preparation of a medicament useful for the treatment of tumors.
12. Use according to claim 11, wherein said tumor is selected from the group consisting of non-small cell lung tumour, tumors of the colon-rectum, prostate, gliomas.
13. Use of a compound of claims 1-4, for the preparation of a medicament useful for the treatment of viral infections.
14. Use of a compound of claims 1-4, for the preparation of a medicament having antiplasmodium falciparum activity.

INTERNATIONAL SEARCH REPORT

Int'l Application No
PCT/IT2004/000118

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D491/22 A61K31/437 A61P35/00
//(C07D491/22,311:00,221:00,221:00,209:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data, PAJ, BEILSTEIN Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 1 044 977 A (IST NAZ STUD CURA DEI TUMORI ; SIGMA TAU IND FARMACEUTI (IT)) 18 October 2000 (2000-10-18) cited in the application claims 1,3; examples CPT154,CPT160 -----	1-14
X	DALLAVALLE, S. ET AL: "Novel cytotoxic 7-iminomethyl and 7-aminomethyl derivatives of camptothecin" BIOORGANIC & MEDICINAL CHEMISTRY LETTERS , 11(3), 291-294 CODEN: BMCLE8; ISSN: 0960-894X, 2001, XP002286415 examples 2b,2f; tables 1,2 ----- -/-	1-14

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

29 June 2004

Date of mailing of the international search report

14/07/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Wörth, C

INTERNATIONAL SEARCH REPORT

In: International Application No

PCT/IT2004/000118

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 97/31003 A (IST NAZ STUD CURA DEI TUMORI ; MERLINI LUCIO (IT); PENCO SERGIO (IT);) 28 August 1997 (1997-08-28) cited in the application claim 1	1-14
Y	EP 0 056 692 A (YAKULT HONSHA KK) 28 July 1982 (1982-07-28) claims 1,11	1-14
Y	EP 0 088 642 A (YAKULT HONSHA KK) 14 September 1983 (1983-09-14) claim 1	1-14
A	SINHA B K: "Topoisomerase Inhibitors. A Review of their Therapeutic Potential in Cancer" DRUGS, ADIS INTERNATIONAL LTD, AT, vol. 49, no. 1, 1995, pages 11-19, XP002116367 ISSN: 0012-6667 the whole document	1-14
A	HUI-KANG WANG ET AL: BIOORGANIC & MEDICINAL CHEMISTRY, ELSEVIER SCIENCE LTD, GB, vol. 2, no. 12, 1994, pages 1397-1402, XP002116366 ISSN: 0968-0896 figure 3; table 4	1-14
A	SAWADA S ET AL: "CHEMICAL MODIFICATION OF AN ANTITUMOR ALKALOID CAMPTOTHECIN: SYNTHESIS AND ANTITUMOR ACTIVITY OF 7-C-SUBSTITUTED CAMPTOTHECINS" CHEMICAL AND PHARMACEUTICAL BULLETIN, PHARMACEUTICAL SOCIETY OF JAPAN. TOKYO, JP, vol. 39, no. 10, 1 October 1991 (1991-10-01), pages 2574-2580, XP002034620 ISSN: 0009-2363 Chart I; table IV	1-14

INTERNATIONAL SEARCH REPORT
Information on patent family members

International Application No
PCT/IT2004/000118

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 1044977	A	18-10-2000	EP 1044977 A1 18-10-2000
			AT 216998 T 15-05-2002
			AU 3160400 A 28-09-2000
			BG 105810 A 28-06-2002
			BR 0008840 A 08-01-2002
			CA 2362760 A1 14-09-2000
			CN 1139592 C 25-02-2004
			CZ 20013077 A3 16-01-2002
			DE 69901379 D1 06-06-2002
			DE 69901379 T2 07-11-2002
			DK 1044977 T3 08-07-2002
			EA 3605 B1 26-06-2003
			EE 200100466 A 16-12-2002
			WO 0053607 A1 14-09-2000
			ES 2175919 T3 16-11-2002
			HK 1031222 A1 09-05-2003
			HR 20010667 A1 31-08-2002
			HU 0200210 A2 29-05-2002
			JP 2002539128 T 19-11-2002
			NO 20014128 A 24-08-2001
			NZ 513393 A 29-04-2003
			PL 355094 A1 22-03-2004
			PT 1044977 T 30-09-2002
			SI 1044977 T1 31-08-2002
			SK 11642001 A3 07-01-2002
			TR 200102603 T2 21-01-2002
			US 2004018988 A1 29-01-2004
			US 6242457 B1 05-06-2001
			US 2001008939 A1 19-07-2001
			ZA 200107408 A 12-03-2002
WO 9731003	A	28-08-1997	IT MI960338 A1 25-08-1997
			AT 221071 T 15-08-2002
			AU 718361 B2 13-04-2000
			AU 1875397 A 10-09-1997
			BR 9707622 A 27-07-1999
			CA 2247513 A1 28-08-1997
			CN 1211987 A ,B 24-03-1999
			CZ 9802652 A3 13-01-1999
			DE 69714220 D1 29-08-2002
			DE 69714220 T2 14-11-2002
			DK 885230 T3 18-11-2002
			EE 9800259 A 15-02-1999
			WO 9731003 A1 28-08-1997
			EP 0885230 A1 23-12-1998
			ES 2180932 T3 16-02-2003
			HU 9900987 A2 30-08-1999
			IL 125856 A 26-08-2001
			JP 2000505448 T 09-05-2000
			NZ 331126 A 28-02-2000
			PT 885230 T 31-12-2002
			RU 2165935 C2 27-04-2001
			SK 112798 A3 11-01-1999
			US 6130227 A 10-10-2000
			US 6306868 B1 23-10-2001
EP 0056692	A	28-07-1982	JP 1438941 C 19-05-1988
			JP 57116075 A 19-07-1982

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int

onal Application No

PCT/IT2004/000118

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0056692	A	JP 62047191 B	06-10-1987
		JP 1438942 C	19-05-1988
		JP 57116076 A	19-07-1982
		JP 62047192 B	06-10-1987
		JP 1438945 C	19-05-1988
		JP 57185285 A	15-11-1982
		JP 62047189 B	06-10-1987
		CA 1177487 A1	06-11-1984
		DE 3265308 D1	19-09-1985
		EP 0056692 A1	28-07-1982
		US 4399276 A	16-08-1983
EP 0088642	A	JP 58154582 A	14-09-1983
	14-09-1983	EP 0088642 A2	14-09-1983